— ABLYBIO, Help Your Research

CD11c Rabbit mAb

货号: AAY1703

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF IP ELISA
推荐浓度	**IHC:** 1:100-1:800 **WB:** 1:1000-1:3000 **IF:** 1:100-1:800 **ELISA:** 1:5000-1:20000 **IP:** 1:50-1:200
理论分子量	128kD
实测分子量	145kD
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.705.
偶联物	Unconjugated
阳性对照	Monocyte,Granulocyte
细胞定位	Membrane>Single-pass type I membrane protein. Cell membrane. Secreted.
纯化	亲和纯化

抗原信息

抗原信息	

靶点信息

研究背景	IL-1R1 (IL-1 receptor type 1, also known as p80, IL-1R-alpha or CD121a) is a 80 kDa transmembrane glycoprotein and a member of the interleukin-1 receptor family (PMID: 35163653; 29248002). It is a receptor for IL-1α, IL-1β, and interleukin-1 receptor antagonist (IL-1Ra). IL-1α and IL-1β interact with the extracellular domain of IL-1R1, triggering the recruitment of an accessory receptor, the IL-1RAcP, resulting in a functional receptor complex that initiates IL-1R1 signaling cascades (PMID: 35163653). IL-1Ra competes with active IL-1 and blocks binding to their common activating receptor IL-1R1 and, thus, inhibits IL-1 signaling (PMID: 35332327; 11641511). It has been reported that IL-1R1 may also bind IL-38 (PMID: 29248002). IL-1R1 expression has been found in many cell types including T cells, macrophages, neutrophils, eosinophils, basophils, mast cells, fibroblasts, and endothelial cells. IL-1R1 can be released from the cell surface following proteolytic cleavage, generating a soluble form of 55-60 kDa (PMID: 17324958).
基因ID	3554;
基因名	ITGAX CD11C
Swiss	P20702 (https://www.uniprot.org/uniprotkb/P20702/entry)
别名	ITGAX CD11C,Integrin alpha-X (CD11 antigen-like family member C) (Leu M5) (Leukocyte adhesion glycoprotein p150,95 alpha chain) (Leukocyte adhesion receptor p150,95) (CD antigen CD11c),Integrin alpha-X, CD11 antigen-like family member C,Leu M5,Leukocyte adhesion glycoprotein p150,95 alpha chain,Leukocyte adhesion receptor p150,95,CD antigen CD11c,Leukocyte ad

产品验证



Western blot analysis of extracts from Mouse thymus tissue using CD11c at 1:1000.



Immunohistochemical analysis of paraffin-embedded human spleen, using CD11c Antibody.，1：200



Immunohistochemical analysis of paraffin-embedded human Lymph node, using CD11c Antibody.，1：200



Immunofluorescence analysis of RAW264.7 cells（LPS stimulate） labelling CD11c with 1：100

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)