— ABLYBIO, Help Your Research

# Bcr Rabbit mAb

**货号: AAY1902**

## 产品信息

| | |
|---|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF IP ELISA |
| 推荐浓度 | **IHC:** 1:100-1:800<br>**WB:** 1:1000-1:3000<br>**IF:** 1:200-1:800<br>**ELISA:** 1:5000-1:20000<br>**IP:** 1:50-1:200 |
| 理论分子量 | 143kD |
| 实测分子量 | 250kD |
| 形式 | Liquid |
| 保存条件 | Store at -20℃. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.75% BSA,50% glycerol,pH7.904. |
| 偶联物 | Unconjugated |
| 阳性对照 | 293T,Jurkat,HeLa,Mouse brain,Rat brain |
| 细胞定位 | Postsynaptic density, Cell projection, dendritic spine, axon, Synapse |
| 纯化 | 亲和纯化 |

## 抗原信息

| | |
|---|---|
| 抗原信息 | |

## 靶点信息

| | |
|---|---|
| 研究背景 | A reciprocal translocation between chromosomes 22 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The protein has serine/threonine kinase activity and is a GTPase-activating protein for p21rac. Two transcript variants encoding different isoforms have been found for this gene. |
| 基因ID | 613 |
| 基因名 | BCR |
| Swiss | P11274 (https://www.uniprot.org/uniprotkb/P11274/entry) |
| 别名 | BCR;Breakpoint cluster region protein;BCR;BCR1;D22S11;Breakpoint cluster region protein;Renal carcinoma antigen NY-REN-26 |

**产品验证**

**实验步骤**

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)