— ABLYBIO, Help Your Research

SQSTM1/p62 (Phospho Ser349) Rabbit mAb

货号: AAY1934

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF ELISA
推荐浓度	**IHC:** 1:100-1:400 **WB:** 1:1000-1:3000 **IF:** 1:100-1:800 **ELISA:** 1:5000-1:20000
理论分子量	48kD
实测分子量	62kD
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.936.
偶联物	Unconjugated
阳性对照	293T,Jurkat
细胞定位	Cytoplasmic vesicle, autophagosome, Preautophagosomal structure, Cytoplasm, cytosol, Nucleus, PML body, Late endosome, Lysosome, Endoplasmic reticulum, myofibril, sarcomere
纯化	亲和纯化

抗原信息

抗原信息	

靶点信息

研究背景	This gene encodes a multifunctional protein that binds ubiquitin and regulates activation of the nuclear fa ctor kappa-B (NF-kB) signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF-kB in response to upstream signals. Alt ernatively spliced transcript variants encoding either the same or different isoforms have been identified for this gene. Mutations in this gene result in sporadic and familial Paget disease of bone.
基因ID	8878
基因名	SQSTM1 ORCA OSIL;SQSTM1/p62 (Ser349)
Swiss	Q13501 (https://www.uniprot.org/uniprotkb/Q13501/entry)
别名	SQSTM1 ORCA OSIL;SQSTM1/p62 (Ser349);Sequestosome-1;EBI3-associated protein of 60 kDa;EBIAP;p60; Phosphotyrosine-independent ligand for the Lck SH2 domain of 62 kDa;Ubiquitin-binding protein p62;

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)