— ABLYBIO, Help Your Research

SFPQ Rabbit mAb

货号: AAY1992

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF ELISA
推荐浓度	**IHC:** 1:200-1:1000 **WB:** 1:2000-1:10000 **IF:** 1:200-1:1000 **ELISA:** 1:5000-1:20000
理论分子量	76kD
实测分子量	100kD
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.994.
偶联物	Unconjugated
阳性对照	OVCAR3,HeLa,Jurkat,HepG2,Mouse brain,Mouse liver,Mouse spleen,Rat brain
细胞定位	Nucleus speckle, Nucleus matrix, Cytoplasm
纯化	亲和纯化

抗原信息

抗原信息	

靶点信息

研究背景	DNA- and RNA binding protein, involved in several nuclear processes. Essential pre-mRNA splicing factor required early in spliceosome formation and for splicing catalytic step II, probably as a heteromer with NONO. Binds to pre-mRNA in spliceosome C complex, and specifically binds to intronic polypyrimidine tracts. Involved in regulation of signal-induced alternative splicing.
基因ID	6421
基因名	SFPQ PSF;Splicing factor, proline- and glutamine-rich (100 kDa DNA-pairing protein) (hPOMp100) (DNA-binding p52/p100 complex, 100 kDa subunit) (Polypyrimidine tract-binding protein-associated-splicing factor) (PSF) (PTB-associated-splicing factor);
Swiss	P23246 (https://www.uniprot.org/uniprotkb/P23246/entry)
别名	SFPQ PSF;Splicing factor, proline- and glutamine-rich (100 kDa DNA-pairing protein) (hPOMp100) (DNA-binding p52/p100 complex, 100 kDa subunit) (Polypyrimidine tract-binding protein-associated-splicing factor) (PSF) (PTB-associated-splicing factor);

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)