— ABLYBIO, Help Your Research

TRIF Rabbit mAb

货号: AAY2024

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF IP ELISA
推荐浓度	**WB:** 1:500-1:2000 **IF:** 1:100-1:800 **ELISA:** 1:5000-1:20000 **IP:** 1:50-1:200
理论分子量	76kD
实测分子量	98kD
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.1026.
偶联物	Unconjugated
阳性对照	Raji,Mouse spleen,Mouse eye,Rat spleen
细胞定位	Cytoplasmic vesicle, autophagosome, Cytoplasm, cytosol, Mitochondrion
纯化	亲和纯化

抗原信息

抗原信息	

靶点信息

研究背景	This gene encodes an adaptor protein containing a Toll/interleukin-1 receptor (TIR) homology domain, which is an intracellular signaling domain that mediates protein-protein interactions between the Toll-like receptors (TLRs) and signal-transduction components. This protein is involved in native immunity against invading pathogens. It specifically interacts with toll-like receptor 3, but not with other TLRs, and this association mediates dsRNA induction of interferon-beta through activation of nuclear factor kappa-B, during an antiviral immune response.
基因ID	148022
基因名	TICAM1
Swiss	Q8IUC6 (https://www.uniprot.org/uniprotkb/Q8IUC6/entry)
别名	TICAM1,TRIF,TIR domain-containing adapter molecule 1,TICAM-1,Proline-rich,vinculin and TIR domain-containing protein B,Putative NF-kappa-B-activating protein 502H,Toll-interleukin-1 receptor domain-containing adapter protein inducing interferon beta,TIR domain-containing adapter protein inducing IFN-beta,TIR

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)