— ABLYBIO, Help Your Research

MED15 Rabbit mAb

货号: AAY2139

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF ELISA
推荐浓度	**IHC:** 1:500-1:2000 **WB:** 1:2000-1:10000 **IF:** 1:200-1:1000 **ELISA:** 1:5000-1:20000
理论分子量	87kD
实测分子量	97kD
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.1141.
偶联物	Unconjugated
阳性对照	Mouse heart,Mouse brain,Rat heart,Rat brain
细胞定位	Cytoplasm, Nucleus
纯化	亲和纯化

抗原信息

抗原信息	

靶点信息

研究背景	The protein encoded by this gene is a subunit of the multiprotein complexes PC2 and ARC/DRIP and may function as a transcriptional coactivator in RNA polymerase II transcription. This gene contains stretches of trinucleotide repeats and is located in the chromosome 22 region which is deleted in DiGeorge syndrome. Alternative splicing results in multiple transcript variants.
基因ID	51586
基因名	MED15 ARC105 CTG7A PCQAP TIG1 TNRC7;Mediator of RNA polymerase II transcription subunit 15 (Activator-recruited cofactor 105 kDa component) (ARC105) (CTG repeat protein 7a) (Mediator complex subunit 15) (Positive cofactor 2 glutamine/Q-rich-associated protein) (PC2 glutamine/Q-rich-associated protein) (TPA-inducible gene 1 protein) (TIG-1) (Trinucleotide repeat-containing gene 7 protein);
Swiss	Q96RN5 (https://www.uniprot.org/uniprotkb/Q96RN5/entry)
别名	MED15 ARC105 CTG7A PCQAP TIG1 TNRC7,Mediator of RNA polymerase II transcription subunit 15 (Activator-recruited cofactor 105 kDa component) (ARC105) (CTG repeat protein 7a) (Mediator complex subunit 15) (Positive cofactor 2 glutamine/Q-rich-associated protein) (PC2 glutamine/Q-rich-associated protein) (TPA-inducible gene 1 protein) (TIG-1) (Trinucleotide repeat-containing gene 7 protein),Mediator of RNA polymerase II transcription subunit 15 (Activator-recruited cofactor 105 kDa component) (ARC105) (CTG repeat protein 7a) (Mediator complex subunit 15) (Positive cofactor 2 glutamine/Q-rich-associated pro

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)