

— ABLYBIO, Help Your Research



# CIITA Antibody

货号: **ABY1023**

## 产品信息

反应	Human, Mouse, Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
理论分子量	123kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.4.
偶联物	Unconjugated
阳性对照	K-562, rat brain
细胞定位	Nucleus
纯化	亲和纯化

## 抗原信息

抗原信息	
------	--

## 靶点信息

研究背景	This gene encodes a protein with an acidic transcriptional activation domain, 4 LRRs (leucine-rich repeats) and a GTP binding domain. The protein is located in the nucleus and acts as a positive regulator of class II major histocompatibility complex gene transcription, and is referred to as the "master control factor" for the expression of these genes. The protein also binds GTP and uses GTP binding to facilitate its own transport into the nucleus. Once in the nucleus it does not bind DNA but rather uses an intrinsic acetyltransferase (AT) activity to act in a coactivator-like fashion. Mutations in this gene have been associated with bare lymphocyte syndrome type II (also known as hereditary MHC class II deficiency or HLA class II-deficient combined immunodeficiency), increased susceptibility to rheumatoid arthritis, multiple sclerosis, and possibly myocardial infarction. Several transcript variants encoding different isoforms have been found for this gene.
基因ID	4261
基因名	CIITA
Swiss	P33076
别名	C2TA; C2TA_HUMAN; CIITA; CIITA IV; Class II major histocompatibility complex transactivator; Class II transactivator; MHC class II transactivator; MHC class II transactivator type III; MHC2TA; NLR family acid domain containing; NLRA; Nucleotide binding oligomerization domain leucine rich repeat and acid domain containing;

## 产品验证

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)