— ABLYBIO, Help Your Research

Phospho-beta Catenin (Ser33/Ser37/Thr41) pAb

货号: ABY1167

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB ICC/IF
推荐浓度	**WB:** 1:500-1:2000,, IF/ICC 1:100-1:500
理论分子量	85kDa
实测分子量	85kDa-92KD
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.5.
偶联物	Unconjugated
阳性对照	293T,Mouse brain,Mouse lung,C6
细胞定位	Cytoplasm
纯化	亲和纯化


抗原信息

抗原信息	


靶点信息

研究背景	The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatrixoma (PTR), medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript variants.
基因ID	1499
基因名	CTNNB1
Swiss	P35222 (https://www.uniprot.org/uniprotkb/P35222/entry)
别名	Beta catenin; Beta-catenin; Cadherin associated protein; Catenin (cadherin associated protein), beta 1, 88kDa; Catenin beta 1; Catenin beta-1; CATNB; CHBCAT; CTNB1_HUMAN; CTNNB; CTNNB1; DKFZp686D02253; FLJ25606; FLJ37923; OTTHUMP00000162082; OTTHUMP00000165222; OTTHUMP00000165223; OTTHUMP00000209288; OTTHUMP00000209289;

产品验证



Western blot analysis of P-beta Catenin (Ser33/Ser37/Thr41) expressed in SW480,HCT-116 ,using P-beta Catenin (Ser33/Ser37/Thr41) Rabbit at 1:500. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s



Immunofluorescence analysis of A549 cells labelling Phospho-beta Catenin (Ser33/Ser37/Thr41) with . The cells were fixed with cold 100% methanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with Goat Anti-Rabbit IgG Fluor488 (1:200，shown in green) at room temprature for 1h.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)