

Mouse anti His-Tag mAb

货号: **AYC25115**

产品信息

反应	All
宿主	Mouse
克隆性	Monoclonal
预测反应	<p>WB: Danio rerio , Homo sapiens , Arabidopsis thaliana , Mus musculus , Other , Saccharomyces cerevisiae , Solanum tuberosum , Caenorhabditis elegans , Crassostrea gigas , Oryza sativa , Aspergillus nidulans , R. anatipestifer , Oryctolagus cuniculus , grass carp , Portunus trituberculatus , Escherichia coli , Nicotiana benthamiana , Cotton bollworms , Cyprinus carpio , Litopenaeus vannamei , Mancinella armigera , Helicoverpa armigera , Sus scrofa , Bos taurus , Ictalurus punctatus , Chlamys Farreri , Zooxanthella , Brachyura , Plasmodium , S. aureus , D. folliculorum and D. brevis adults</p> <p>IF: Homo sapiens , Saccharomyces cerevisiae , grass carp , Rattus norvegicus , Mus musculus , Danio rerio , Cyprinus carpio</p> <p>IP: Homo sapiens , Cyprinus carpio , Mus musculus , 无 , Rattus norvegicus</p> <p>ELISA: Cynoglossus semilaevis , Mus musculus , Rattus norvegicus , Oyster</p> <p>Co-IP: Helicoverpa armigera , Homo sapiens</p> <p>CoIP: B. napus</p> <p>co-ip: Homo sapiens</p> <p>IF、IHC: Mus musculus</p> <p>PAMPs binding assay: OystersC. gigas</p> <p>IHC: Homo sapiens</p>
应用	WB IP
推荐浓度	<p>WB: 1:2000 - 1:20000</p> <p>IP: 1:50 - 1:100</p>
理论分子量	0.8KDa
实测分子量	27KDa,35KDa
形式	Liquid
保存条件	<p>Store at -20°C. Avoid freeze / thaw cycles.</p> <p>Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.</p>
偶联物	Unconjugated
阳性对照	293T
细胞定位	
纯化	Affinity purification

抗原信息

抗原信息	A synthetic peptide corresponding to His tag.
------	---

靶点信息

研究背景	Consecutive histidine residues (usually 6 to 10 in length) are often inserted into the amino acid sequences of recombinant proteins. The resulting His-tagged proteins can be detected by using His antibodies.
基因ID	
基因名	
Swiss	
别名	His;His tag;His-tag

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn