— ABLYBIO, Help Your Research

Mouse anti HA-Tag mAb

货号: AYC25116

产品信息

反应	All
宿主	Mouse
克隆性	Monoclonal
预测反应	**WB:** Arabidopsis thaliana , Mus musculus , Drosophila melanogaster , Homo sapiens , Solanum tuberosum L. , M. oryzae , Sus scrofa , Serpentiformes , Penicillium oxalicum , Nicotiana tabacum L , Mycobacterium tuberculosis , Other **Co-IP:** Oryza sativa , Arabidopsis thaliana , Mus musculus , Homo sapiens , Cucumis sativus , Sac fungi , rainbow trout **IP:** Other , Nicotiana tabacum , Arabidopsis , Homo sapiens , Mus musculus , Vitis vinifera , Sus scrofa **Yeast three-hybrid assay:** Oryza sativa **IHC:** Mus musculus,Homo sapiens **GPCRs:** Mus musculus **IF:** Homo sapiens,Serpentiformes , Sus scrofa , Cyprinus gibelio , Mus musculus **CoIP:** Arabidopsis thaliana **co-IP:** Mus musculus **ELISA:** Ctenopharyngodon idellus **CO-IP:** Homo sapiens
应用	WB IP ICC/IF
推荐浓度	**WB:** 1:2000 - 1:20000 **IF/ICC:** 1:100 - 1:500 **IP:** 1:20 - 1:50
理论分子量	
实测分子量	
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to HA tag.

靶点信息

研究背景	Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes.Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.
基因ID	
基因名	
Swiss	
别名	HA,HA tag,HA-tag,Mouse anti HA-Tag mAb

产品验证



Western blot analysis of HA-Tag expressed in 293T using Rabbit anti HA-Tag pAb at 1:2000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)