

— ABLYBIO, Help Your Research



SH2D1A/SAP (YD13397) Rabbit mAb

货号: **AYD11045**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	
理论分子量	14kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,PC-3,Mouse spleen
细胞定位	Cytoplasm
纯化	亲和纯化



抗原信息

抗原信息	
------	--

靶点信息

研究背景	This gene encodes a protein that plays a major role in the bidirectional stimulation of T and B cells. This protein contains an SH2 domain and a short tail. It associates with the signaling lymphocyte-activation molecule, thereby acting as an inhibitor of this transmembrane protein by blocking the recruitment of the SH2-domain-containing signal-transduction molecule SHP-2 to its docking site. This protein can also bind to other related surface molecules that are expressed on activated T, B and NK cells, thereby modifying signal transduction pathways in these cells. Mutations in this gene cause lymphoproliferative syndrome X-linked type 1 or Duncan disease, a rare immunodeficiency characterized by extreme susceptibility to infection with Epstein-Barr virus, with symptoms including severe mononucleosis and malignant lymphoma. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	4068
基因名	SH2D1A
Swiss	O60880 (https://www.uniprot.org/uniprotkb/O60880/entry)
别名	SH2D1A/SAP (YD13397),SH2D1A/SAP (YD13397) Rabbit mAb,SH2D1A,Duncan disease SH2-protein,Signaling lymphocytic activation molecule-associated protein,T-cell signal transduction molecule SAP,DSHP,SAP

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)