

— ABLYBIO, Help Your Research



# Kappa Light chain (YD20875) Rabbit mAb

货号: **AYD11135**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P IP
推荐浓度	
理论分子量	12kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Human serum,Mouse serum,Rat serum
细胞定位	Secreted, Cell membrane
纯化	亲和纯化


## 抗原信息

抗原信息	
------	--

## 靶点信息

研究背景	Constant region of immunoglobulin light chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulin-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens. The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D-J) rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen.
基因ID	3514
基因名	IGKC
Swiss	P01834 ( <a href="https://www.uniprot.org/uniprotkb/P01834/entry">https://www.uniprot.org/uniprotkb/P01834/entry</a> )
别名	Kappa Light chain (YD20875),Kappa Light chain (YD20875) Rabbit mAb,IGKC,Ig kappa chain C region,Ig kappa chain C region AG,Ig kappa chain C region CUM,Ig kappa chain C region EU,Ig kappa chain C region OU,Ig kappa chain C region ROY,Ig kappa chain C region TI

## 产品验证



## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)