

Bag3 (YD19448) Rabbit mAb

货号: **AYD11155**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB FC |
| 推荐浓度 | |
| 理论分子量 | 62kDa |
| 实测分子量 | |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | A-549,HeLa,HepG2,Mouse brain,Mouse testis |
| 细胞定位 | Nucleus, Cytoplasm |
| 纯化 | |

抗原信息


| | |
|------|--|
| 抗原信息 | |
|------|--|

靶点信息

| | |
|------|---|
| 研究背景 | BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. |
| 基因ID | 9531 |
| 基因名 | BAG3 |

| | |
|-------|----------------|
| Swiss | O95817 |
| 别名 | Bag3 (YD19448) |

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn