

— ABLYBIO, Help Your Research



USP16 (YD33322) Rabbit mAb

货号: **AYD11574**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	
理论分子量	93kDa/94kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Nucleus, Cytoplasm
纯化	亲和纯化


抗原信息

抗原信息	
------	--

靶点信息

研究背景	Specifically deubiquitinates 'Lys-120' of histone H2A (H2AK119Ub), a specific tag for epigenetic transcriptional repression, thereby acting as a coactivator (PubMed:24034696). Deubiquitination of histone H2A is a prerequisite for subsequent phosphorylation at 'Ser-11' of histone H3 (H3S10ph), and is required for chromosome segregation when cells enter into mitosis (PubMed:24034696). In resting B- and T-lymphocytes, phosphorylation by AURKB leads to enhance its activity, thereby maintaining transcription in resting lymphocytes (PubMed:24034696). Regulates Hox gene expression via histone H2A deubiquitination. Prefers nucleosomal substrates. Does not deubiquitinate histone H2B. Also deubiquitinates non-histone proteins, such as ribosomal protein RPS27A: deubiquitination of monoubiquitinated RPS27A promotes maturation of the 40S ribosomal subunit. Also mediates deubiquitination of tektin proteins (TEKT1, TEKT2, TEKT3, TEKT4 and TEKT5), promoting their stability (PubMed:36206347) Specifically deubiquitinates 'Lys-120' of histone H2A (H2AK119Ub), a specific tag for epigenetic transcriptional repression, thereby acting as a coactivator. Deubiquitination of histone H2A is a prerequisite for subsequent phosphorylation at 'Ser-11' of histone H3 (H3S10ph), and is required for chromosome segregation when cells enter into mitosis. In resting B- and T-lymphocytes, phosphorylation by AURKB leads to enhance its activity, thereby maintaining transcription in resting lymphocytes. Regulates Hox gene expression via histone H2A deubiquitination. Prefers nucleosomal substrates. Does not deubiquitinate histone H2B. Also deubiquitinates non-histone proteins, such as ribosomal protein RPS27A: deubiquitination of monoubiquitinated RPS27A promotes maturation of the 40S ribosomal subunit. Also mediates deubiquitination of tektin proteins (TEKT1, TEKT2, TEKT3, TEKT4 and TEKT5), promoting their stability
基因ID	3134
基因名	Usp16
Swiss	Q99LG0 (https://www.uniprot.org/uniprotkb/Q99LG0/entry), Q2KJ09 (https://www.uniprot.org/uniprotkb/Q2KJ09/entry)
别名	USP16 (YD33322),USP16 (YD33322) Rabbit mAb,Usp16,Deubiquitinating enzyme 16,Ubiquitin thioesterase 16,Ubiquitin-specific-processing protease 16

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)