

— ABLYBIO, Help Your Research



Eif4a3 (YD13806) Rabbit mAb

货号: **AYD11948**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P ICC/IF FC IP
推荐浓度	
理论分子量	47kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,Jurkat,HeLa,A-549,NIH/3T3,Mouse spleen,Rat kidney
细胞定位	Nucleus, Nucleus speckle, Cytoplasm
纯化	亲和纯化




抗原信息

抗原信息	
------	--

靶点信息

研究背景	This gene encodes a member of the DEAD box protein family. DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure, such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. The protein encoded by this gene is a nuclear matrix protein. Its amino acid sequence is highly similar to the amino acid sequences of the translation initiation factors eIF4AI and eIF4AII, two other members of the DEAD box protein family.
基因ID	9775
基因名	EIF4A3
Swiss	P38919 (https://www.uniprot.org/uniprotkb/P38919/entry)
别名	Eif4a3 (YD13806), Eif4a3 (YD13806) Rabbit mAb, EIF4A3, ATP-dependent RNA helicase DDX48, ATP-dependent RNA helicase eIF4A-3, DEAD box protein 48, Eukaryotic initiation factor 4A-like NUK-34, Eukaryotic translation initiation factor 4A isoform 3, Nuclear matrix protein 265, DDX48

产品验证



实验步骤

