

— ABLYBIO, Help Your Research



NEDD8 Activating Enzyme E1C (YD33652) Rabbit mAb

货号: **AYD12248**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	
理论分子量	52kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,HepG2,K-562,RAW264.7,Rat heart
细胞定位	
纯化	亲和纯化



抗原信息

抗原信息	
------	--

靶点信息

研究背景	The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E1 ubiquitin-activating enzyme family. The encoded enzyme associates with AppBp1, an amyloid beta precursor protein binding protein, to form a heterodimer, and then the enzyme complex activates NEDD8, a ubiquitin-like protein, which regulates cell division, signaling and embryogenesis. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.
基因ID	9039
基因名	UBA3
Swiss	Q8TBC4 (https://www.uniprot.org/uniprotkb/Q8TBC4/entry)
别名	NEDD8 Activating Enzyme E1C (YD33652),NEDD8 Activating Enzyme E1C (YD33652) Rabbit mAb,UBA3,NEDD8-activating enzyme E1C,Ubiquitin-activating enzyme E1C,Ubiquitin-like modifier-activating enzyme 3,UBE1C

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)