

— ABLYBIO, Help Your Research



uPA Receptor/U-PAR (YD16933) Rabbit mAb

货号: **AYD12908**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	ICC/IF FC IP
推荐浓度	
理论分子量	37kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	DU145,HT-1080,A-549,HeLa,Mouse lung,Rat thymus
细胞定位	Cell membrane, Cell projection, invadopodium membrane, Secreted
纯化	亲和纯化

抗原信息

抗原信息	
------	--

靶点信息

研究背景	This gene encodes the receptor for urokinase plasminogen activator and, given its role in localizing and promoting plasmin formation, likely influences many normal and pathological processes related to cell-surface plasminogen activation and localized degradation of the extracellular matrix. It binds both the proprotein and mature forms of urokinase plasminogen activator and permits the activation of the receptor-bound pro-enzyme by plasmin. The protein lacks transmembrane or cytoplasmic domains and may be anchored to the plasma membrane by a glycosyl-phosphatidylinositol (GPI) moiety following cleavage of the nascent polypeptide near its carboxy-terminus. However, a soluble protein is also produced in some cell types. Alternative splicing results in multiple transcript variants encoding different isoforms. The proprotein experiences several post-translational cleavage reactions that have not yet been fully defined.
基因ID	5329
基因名	PLAUR
Swiss	Q03405 (https://www.uniprot.org/uniprotkb/Q03405/entry)
别名	uPA Receptor/U-PAR (YD16933),uPA Receptor/U-PAR (YD16933) Rabbit mAb,PLAUR,Monocyte activation antigen Mo3,MO3,UPAR

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)