

— ABLYBIO, Help Your Research



SIRT2 (YD12954) Rabbit mAb

货号: **AYD13593**

产品信息

| | |
|-------|---|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | |
| 理论分子量 | 43kDa |
| 实测分子量 | |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse brain, Rat brain |
| 细胞定位 | Nucleus, Cytoplasm, perinuclear region, cytoskeleton, microtubule organizing center, centrosome, centriole, spindle, Midbody, Chromosome, Perikaryon, Cell projection, growth cone, Myelin membrane |
| 纯化 | 亲和纯化 |


抗原信息

| | |
|------|--|
| 抗原信息 | |
|------|--|

靶点信息

| | |
|-------|--|
| 研究背景 | This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Several transcript variants are resulted from alternative splicing of this gene. |
| 基因ID | 22933 |
| 基因名 | SIRT2 |
| Swiss | Q8IXJ6 (https://www.uniprot.org/uniprotkb/Q8IXJ6/entry) |
| 别名 | SIRT2 (YD12954),SIRT2 (YD12954) Rabbit mAb,SIRT2,NAD-dependent protein deacylase sirtuin-2,NAD-dependent protein defatty-acylase sirtuin-2,Regulatory protein SIR2 homolog 2,SIR2-like protein 2,SIR2L,SIR2L2 |

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)