

— ABLYBIO, Help Your Research



CD86 (YD31778) Rabbit mAb

货号: **AYD13648**

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	FC
推荐浓度	
理论分子量	35kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cell membrane
纯化	亲和纯化

抗原信息

抗原信息	
------	--

靶点信息

研究背景	Costimulatory molecule that belongs to the immunoglobulin superfamily that plays an important role in T-lymphocyte activation (By similarity). Acts as the primary auxiliary signal augmenting the MHC/TCR signal in naive T-cells by acting as a ligand for the CD28 receptor which is constitutively expressed on the cell surface of T-cells (By similarity). May play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation (By similarity). Also involved in the regulation of B cells function, plays a role in regulating the level of IgG(1) produced (PubMed:23241883). Upon CD40 engagement, activates NF-kappa-B signaling pathway via phospholipase C and protein kinase C activation (PubMed:23241883). Also acts as an inhibitor of T-cell activation by acting as a ligand for CTLA4, a decoy receptor, thereby blocking CD28-mediated T-cell priming (PubMed:7504059, PubMed:7513726)
基因ID	3134
基因名	Cd86
Swiss	P42082 (https://www.uniprot.org/uniprotkb/P42082/entry)
别名	CD86 (YD31778),CD86 (YD31778) Rabbit mAb,Cd86,Activation B7-2 antigen,Early T-cell costimulatory molecule 1

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)