

— ABLYBIO, Help Your Research



NUP98 (YD32200) Rabbit mAb

货号: **AYD13882**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P ICC/IF
推荐浓度	
理论分子量	198kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,LO2,HeLa,Jurkat,HL-60
细胞定位	Nucleus membrane, Nucleus, nuclear pore complex, nucleoplasm
纯化	亲和纯化


抗原信息

抗原信息	
------	--

靶点信息

研究背景	Nuclear pore complexes (NPCs) regulate the transport of macromolecules between the nucleus and cytoplasm, and are composed of many polypeptide subunits, many of which belong to the nucleoporin family. This gene belongs to the nucleoporin gene family and encodes a 186 kDa precursor protein that undergoes autoproteolytic cleavage to generate a 98 kDa nucleoporin and 96 kDa nucleoporin. The 98 kDa nucleoporin contains a Gly-Leu-Phe-Gly (GLGF) repeat domain and participates in many cellular processes, including nuclear import, nuclear export, mitotic progression, and regulation of gene expression. The 96 kDa nucleoporin is a scaffold component of the NPC. Proteolytic cleavage is important for targeting of the proteins to the NPC. Translocations between this gene and many other partner genes have been observed in different leukemias. Rearrangements typically result in chimeras with the N-terminal GLGF domain of this gene to the C-terminus of the partner gene. Alternative splicing results in multiple transcript variants encoding different isoforms, at least two of which are proteolytically processed. Some variants lack the region that encodes the 96 kDa nucleoporin.
基因ID	4928
基因名	NUP98
Swiss	P52948 (https://www.uniprot.org/uniprotkb/P52948/entry)
别名	NUP98 (YD32200),NUP98 (YD32200) Rabbit mAb,NUP98,98 kDa nucleoporin,Nucleoporin Nup98,96 kDa nucleoporin,Nucleoporin Nup96,ADAR2

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)