

Phospho-Bcr (Tyr177) (YD12787) Rabbit mAb

货号: **AYD14074**

产品信息

| | |
|-------|---|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | |
| 理论分子量 | 143kDa |
| 实测分子量 | |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | 293T,Jurkat,HeLa,Mouse brain,Rat brain |
| 细胞定位 | Postsynaptic density, Cell projection, dendritic spine, axon, Synapse |
| 纯化 | |

抗原信息

| | |
|------|--|
| 抗原信息 | |
|------|--|

靶点信息

| | |
|------|---|
| 研究背景 | A reciprocal translocation between chromosomes 22 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The protein has serine/threonine kinase activity and is a GTPase-activating protein for p21rac. Two transcript variants encoding different isoforms have been found for this gene. |
| 基因ID | 613 |
| 基因名 | BCR |

| | |
|--------------|--------------------------------|
| Swiss | P11274 |
| 别名 | Phospho-Bcr (Tyr177) (YD12787) |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn