

EMAP II (YD19900) Rabbit mAb

货号: **AYD14164**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | |
| 理论分子量 | 34kDa |
| 实测分子量 | |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HL-60,22RV1,HepG2,HT-29,MCF7,A-549 |
| 细胞定位 | Nucleus, Cytoplasm, cytosol, Secreted, Endoplasmic reticulum, Golgi apparatus |
| 纯化 | |

抗原信息



| | |
|------|--|
| 抗原信息 | |
|------|--|

靶点信息

| | |
|------|--|
| 研究背景 | The protein encoded by this gene is a cytokine that is specifically induced by apoptosis, and it is involved in the control of angiogenesis, inflammation, and wound healing. The release of this cytokine renders the tumor-associated vasculature sensitive to tumor necrosis factor. The precursor protein is identical to the p43 subunit, which is associated with the multi-tRNA synthetase complex, and it modulates aminoacylation activity of tRNA synthetase in normal cells. This protein is also involved in the stimulation of inflammatory responses after proteolytic cleavage in tumor cells. Multiple transcript variants encoding different isoforms have been found for this gene. A pseudogene has been identified on chromosome 20. |
| 基因ID | 9255 |
| 基因名 | AIMP1 |

| | |
|-------|-------------------|
| Swiss | Q12904 |
| 别名 | EMAP II (YD19900) |

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn