— ABLYBIO, Help Your Research

CD22 (YD19053) Rabbit mAb

货号: AYD14465

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P ICC/IF IP
推荐浓度	
理论分子量	95kDa
实测分子量	
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse thymus,Rat lung
细胞定位	Cell membrane
纯化	亲和纯化


抗原信息


抗原信息	

靶点信息

研究背景	Siglecs (sialic acid binding Ig-like lectins) are I-type (Ig-type) lectins belonging to the Ig superfamily. They are characterized by an N-terminal Ig-like V-type domain which mediates sialic acid binding, followed by v arying numbers of Ig-like C2-type domains. Human Siglec-2, also known as B-cell antigen CD22 or Blymp hocyte cell adhesion molecule (BL-CAM), is a B-cell restricted glycoprotein that is expressed in the cytopl asm of progenitor B and pre-B cells and on the surface of mature B cells. Two distinct human Siglec2/CD2 2 cDNAs that arise from differential RNA processing of the same gene have been isolated. Siglec2/CD22 i s an adhesion molecule that preferentially binds alpha 2,6- linked sialic acid on the same (cis) or adjacent (trans) cells. Interaction of CD22 with trans ligands on opposing cells was found to be favored over the bi nding of ligands in cis.
基因ID	933
基因名	CD22
Swiss	P20273 (https://www.uniprot.org/uniprotkb/P20273/entry)
别名	CD22 (YD19053),CD22 (YD19053) Rabbit mAb,CD22,B-lymphocyte cell adhesion molecule,Sialic acid-bin ding Ig-like lectin 2,T-cell surface antigen Leu-14,SIGLEC2

产品验证





实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)