

— ABLYBIO, Help Your Research



# FGF2 (YD19041) Rabbit mAb

货号: **AYD14571**

## 产品信息

|       |   |
|-------|---|
| 反应    | Human   |
| 宿主    | Rabbit  |
| 克隆性   | Monoclonal  |
| 预测反应  |   |
| 应用    | WB IHC-P IP   |
| 推荐浓度  |   |
| 理论分子量 | 31kDa   |
| 实测分子量 |   |
| 形式    | Liquid  |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated  |
| 阳性对照  | 293T  |
| 细胞定位  | Secreted, Nucleus   |
| 纯化    | 亲和纯化  |


## 抗原信息

|      |  |
|------|--|
| 抗原信息 |  |
|------|--|

## 靶点信息

|       |   |
|-------|---|
| 研究背景  | The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF. |
| 基因ID  | 2247  |
| 基因名   | FGF2  |
| Swiss | P09038 ( <a href="https://www.uniprot.org/uniprotkb/P09038/entry">https://www.uniprot.org/uniprotkb/P09038/entry</a> )  |
| 别名    | FGF2 (YD19041),FGF2 (YD19041) Rabbit mAb,FGF2,Basic fibroblast growth factor,Heparin-binding growth factor 2,FGFB   |

## 产品验证



## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)