

XPD (YD16282) Rabbit mAb

货号: **AYD14606**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P ICC/IF
推荐浓度	
理论分子量	87kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1,293T,HepG2
细胞定位	Nucleus, Cytoplasm, cytoskeleton, spindle
纯化	

抗原信息



抗原信息	
------	--

靶点信息

研究背景	The nucleotide excision repair pathway is a mechanism to repair damage to DNA. The protein encoded by this gene is involved in transcription-coupled nucleotide excision repair and is an integral member of the basal transcription factor BTF2/TFIIH complex. The gene product has ATP-dependent DNA helicase activity and belongs to the RAD3/XPD subfamily of helicases. Defects in this gene can result in three different disorders, the cancer-prone syndrome xeroderma pigmentosum complementation group D, trichothiodystrophy, and Cockayne syndrome. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
基因ID	2068
基因名	ERCC2

Swiss	P18074
别名	XPD (YD16282)

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn