

— ABLYBIO, Help Your Research



eIF2A (YD12621) Rabbit mAb

货号: **AYD14669**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P ICC/IF
推荐浓度	
理论分子量	65kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,PC-3,U-251MG,Mouse liver
细胞定位	
纯化	亲和纯化




抗原信息


抗原信息	
------	--

靶点信息

研究背景	This gene encodes a eukaryotic translation initiation factor that catalyzes the formation of puromycin-sensitive 80 S preinitiation complexes and the poly(U)-directed synthesis of polyphenylalanine at low concentrations of Mg ²⁺ . This gene should not be confused with eIF2-alpha (EIF2S1, Gene ID: 1965), the alpha subunit of the eIF2 translation initiation complex. Although both of these proteins function in binding initiator tRNA to the 40 S ribosomal subunit, the encoded protein does so in a codon-dependent manner, whereas eIF2 complex requires GTP. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms.
基因ID	83939
基因名	EIF2A
Swiss	Q9BY44 (https://www.uniprot.org/uniprotkb/Q9BY44/entry)
别名	eIF2A (YD12621),eIF2A (YD12621) Rabbit mAb,EIF2A,65 kDa eukaryotic translation initiation factor 2A

产品验证





实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)