

— ABLYBIO, Help Your Research



# CD80 (YD16488) Rabbit mAb

货号: **AYD14856**

## 产品信息

反应	Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	
理论分子量	35kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cell membrane
纯化	亲和纯化


## 抗原信息

抗原信息	
------	--

## 靶点信息

研究背景	Costimulatory molecule that belongs to the immunoglobulin superfamily that plays an important role in T-lymphocyte activation (PubMed:16339518, PubMed:1714935). Acts as the primary auxiliary signal augmenting the MHC/TCR signal in naive T-cells by acting as a ligand for the CD28 receptor which is constitutively expressed on the cell surface of T-cells (PubMed:16339518). In turn, activates different signaling pathways such as NF-kappa-B or MAPK leading to the production of different cytokines (By similarity). Also acts as an inhibitor of T-cell activation by acting as a ligand for CTLA4, a decoy receptor, thereby blocking CD28-mediated T-cell priming (By similarity). In addition, CD28/CD80 costimulatory signal stimulates glucose metabolism and ATP synthesis of T-cells by activating the PI3K/Akt signaling pathway (By similarity). Also acts as a regulator of PDL1/PDCD1 interactions to limit excess engagement of PDL1 and its inhibitory role in immune responses (By similarity). Expressed on B-cells, plays a critical role in regulating interactions between B-cells and T-cells in both early and late germinal center responses, which are crucial for the generation of effective humoral immune responses (PubMed:22450810)
基因ID	3134
基因名	Cd80
Swiss	Q00609 ( <a href="https://www.uniprot.org/uniprotkb/Q00609/entry">https://www.uniprot.org/uniprotkb/Q00609/entry</a> )
别名	CD80 (YD16488),CD80 (YD16488) Rabbit mAb,Cd80,Activation B7-1 antigen,B7

## 产品验证



## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)