

— ABLYBIO, Help Your Research



Ku70 (YD33395) Rabbit mAb

货号: **AYD14903**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P ICC/IF
推荐浓度	
理论分子量	70kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	COS-7,SW480,Mouse brain
细胞定位	Nucleus, Chromosome, Cytoplasm
纯化	亲和纯化




抗原信息


抗原信息	
------	--

靶点信息

研究背景	The p70/p80 autoantigen is a nuclear complex consisting of two subunits with molecular masses of approximately 70 and 80 kDa. The complex functions as a single-stranded DNA-dependent ATP-dependent helicase. The complex may be involved in the repair of nonhomologous DNA ends such as that required for double-strand break repair, transposition, and V(D)J recombination. High levels of autoantibodies to p70 and p80 have been found in some patients with systemic lupus erythematosus.
基因ID	2547
基因名	XRCC6
Swiss	P12956 (https://www.uniprot.org/uniprotkb/P12956/entry)
别名	Ku70 (YD33395),Ku70 (YD33395) Rabbit mAb,XRCC6,5'-deoxyribose-5-phosphate lyase Ku70,70 kDa subunit of Ku antigen,ATP-dependent DNA helicase 2 subunit 1,ATP-dependent DNA helicase II 70 kDa subunit, CTC box-binding factor 75 kDa subunit,DNA repair protein XRCC6

产品验证





实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)