— ABLYBIO, Help Your Research

Wnt3a (YD15779) Rabbit mAb

货号: AYD14992

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC/IF FC IP
推荐浓度	**WB: 1:1000-1：3000:** **IHC-P: 1:100-1:500:** **ICC/IF: 1:100-1：300:** **FC: 1:200-1:500:** **IP: 1:20-1:50:**
理论分子量	39kDa
实测分子量	39 kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U2OS
细胞定位	Secreted, extracellular space, extracellular matrix
纯化	亲和纯化


抗原信息

抗原信息	


靶点信息

研究背景	The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It encodes a protein which shows 96% amino acid identity to mouse Wnt3A protein, and 84% to human WNT3 protein, another WNT gene product. This gene is clustered with WNT14 gene, another family member, in chromosome 1q42 region.
基因ID	89780
基因名	WNT3A
Swiss	P56704 (https://www.uniprot.org/uniprotkb/P56704/entry)
别名	Wnt3a (YD15779),Wnt3a (YD15779) Rabbit mAb,WNT3A

产品验证



Immunohistochemical analysis of formalin fixed paraffin embedded human Lung adenocarcinoma tissue with Wnt3a Rabbit mAb，1：200



Immunofluorescence analysis of SH-SY5Y cells labelling Wnt3a with .，1：200 The cells were fixed with cold 100% methanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with Goat Anti-Rabbit IgG Fluor488 (1:200，shown in green) at room temprature for 1h.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)