

HDAC6 (YD15259) Rabbit mAb

货号: **AYD15099**

产品信息

| | |
|-------|---|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC-P |
| 推荐浓度 | |
| 理论分子量 | 131kDa |
| 实测分子量 | |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Rat liver |
| 细胞定位 | Cytoplasm, cytoskeleton, Nucleus, Perikaryon, Cell projection, dendrite, axon, cilium, microtubule organizing center, centrosome, cilium basal body |
| 纯化 | |

抗原信息

| | |
|------|--|
| 抗原信息 | |
|------|--|

靶点信息

| | |
|------|---|
| 研究背景 | Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events . Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It contains an internal duplication of two catalytic domains which appear to function independently of each other. This protein possesses histone deacetylase activity and represses transcription. |
| 基因ID | 10013 |
| 基因名 | HDAC6 |

| | |
|--------------|-----------------|
| Swiss | Q9UBN7 |
| 别名 | HDAC6 (YD15259) |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn