

— ABLYBIO, Help Your Research



TGF beta 1 (YD15235) Rabbit mAb

货号: **AYD15107**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC-P |
| 推荐浓度 | |
| 理论分子量 | 44kDa |
| 实测分子量 | |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | K-562,Mouse spleen,Rat lung |
| 细胞定位 | Secreted, extracellular space, extracellular matrix |
| 纯化 | 亲和纯化 |



抗原信息

| | |
|------|--|
| 抗原信息 | |
|------|--|

靶点信息

| | |
|-------|--|
| 研究背景 | This gene encodes a member of the transforming growth factor beta (TGFB) family of cytokines, which are multifunctional peptides that regulate proliferation, differentiation, adhesion, migration, and other functions in many cell types. Many cells have TGFB receptors, and the protein positively and negatively regulates many other growth factors. The secreted protein is cleaved into a latency-associated peptide (LAP) and a mature TGFB1 peptide, and is found in either a latent form composed of a TGFB1 homodimer, a LAP homodimer, and a latent TGFB1-binding protein, or in an active form composed of a TGFB1 homodimer. The mature peptide may also form heterodimers with other TGFB family members. This gene is frequently upregulated in tumor cells, and mutations in this gene result in Camurati-Engelmann disease. |
| 基因ID | 7040 |
| 基因名 | TGFB1 |
| Swiss | P01137 (https://www.uniprot.org/uniprotkb/P01137/entry) |
| 别名 | TGF beta 1 (YD15235),TGF beta 1 (YD15235) Rabbit mAb,TGFB1,TGFB |

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)