

— ABLYBIO, Help Your Research



Granulin (YD15099) Rabbit mAb

货号: **AYD15145**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P ICC/IF FC IP
推荐浓度	
理论分子量	64kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HL-60,THP-1,PC-12,293T,HepG2,MCF7,Mouse liver,Mouse spleen,Mouse kidney
细胞定位	Secreted, Lysosome
纯化	亲和纯化

抗原信息

抗原信息	
------	--

靶点信息

研究背景	Granulins are a family of secreted, glycosylated peptides that are cleaved from a single precursor protein with 7.5 repeats of a highly conserved 12-cysteine granulin/epithelin motif. The 88 kDa precursor protein, progranulin, is also called proepithelin and PC cell-derived growth factor. Cleavage of the signal peptide produces mature granulin which can be further cleaved into a variety of active, 6 kDa peptides. These smaller cleavage products are named granulin A, granulin B, granulin C, etc. Epithelins 1 and 2 are synonymous with granulins A and B, respectively. Both the peptides and intact granulin protein regulate cell growth. However, different members of the granulin protein family may act as inhibitors, stimulators, or have dual actions on cell growth. Granulin family members are important in normal development, wound healing, and tumorigenesis.
基因ID	2896
基因名	GRN
Swiss	P28799 (https://www.uniprot.org/uniprotkb/P28799/entry)
别名	Granulin (YD15099),Granulin (YD15099) Rabbit mAb,GRN,Acrogranin,Epithelin precursor,Glycoprotein of 88 Kda,Granulin precursor,PC cell-derived growth factor,Proepithelin,Granulin G,Granulin F,Epithelin-2,Granulin B,Epithelin-1,Granulin A,Granulin C,Granulin D,Granulin E

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)