

Ogg1 (YD14232) Rabbit mAb

货号: **AYD15377**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	
理论分子量	39kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	Nucleus, nucleoplasm, Nucleus speckle, Nucleus matrix, Mitochondrion
纯化	

抗原信息


抗原信息	
------	--

靶点信息

研究背景	<p>This gene encodes the enzyme responsible for the excision of 8-oxoguanine, a mutagenic base byproduct which occurs as a result of exposure to reactive oxygen. The action of this enzyme includes lyase activity for chain cleavage. Alternative splicing of the C-terminal region of this gene classifies splice variants into two major groups, type 1 and type 2, depending on the last exon of the sequence. Type 1 alternative splice variants end with exon 7 and type 2 end with exon 8. All variants share the N-terminal region in common, which contains a mitochondrial targeting signal that is essential for mitochondrial localization. Many alternative splice variants for this gene have been described, but the full-length nature for every variant has not been determined.</p>
基因ID	4968

基因名	OGG1
Swiss	O15527
别名	Ogg1 (YD14232)

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn