

# CARD12 (YD14227) Rabbit mAb

货号: **AYD15380**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	
理论分子量	117kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cytoplasm, cytosol, Inflammasome
纯化	

## 抗原信息

抗原信息	
------	--


## 靶点信息

研究背景	<p>Key component of inflammasomes that indirectly senses specific proteins from pathogenic bacteria and fungi and responds by assembling an inflammasome complex that promotes caspase-1 activation, cytokine production and macrophage pyroptosis. The NLRC4 inflammasome is activated as part of the innate immune response to a range of intracellular bacteria. It senses pathogenic proteins of the type III secretion system (T3SS) and type IV secretion system (T4SS) such as flagellin and PrgJ-like rod proteins via the Naip proteins (Naip1, Naip2 or Naip5): specific Naip proteins recognize and bind pathogenic proteins, driving assembly and activation of the NLRC4 inflammasome. The NLRC4 inflammasome senses Gram-negative bacteria such as <i>L.pneumophila</i> and <i>P.aeruginosa</i>, enteric pathogens <i>S.typhimurium</i> (<i>Salmonella</i>) and <i>S.fl exneri</i> and fungal pathogen <i>C.albicans</i>. In intestine, the NLRC4 inflammasome is able to discriminate between commensal and pathogenic bacteria and specifically drives production of interleukin-1 beta (IL1B) in response to infection by <i>Salmonella</i> or <i>P.aeruginosa</i>. In case of <i>L.pneumophila</i> infection the inflammasome acts by activating caspase-7</p>
------	--

基因ID	3134
基因名	Nlrc4
Swiss	Q3UP24
别名	CARD12 (YD14227)

## 产品验证

---



## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)