

ACAA2 (YD36015) Rabbit mAb

货号: **AYD16125**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P ICC/IF
推荐浓度	
理论分子量	42kDa/42kDa/42kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Mitochondrion
纯化	



抗原信息

抗原信息	
------	--

靶点信息

研究背景	In the production of energy from fats, this is one of the enzymes that catalyzes the last step of the mitochondrial beta-oxidation pathway, an aerobic process breaking down fatty acids into acetyl-CoA (Probable). Using free coenzyme A/CoA, catalyzes the thiolitic cleavage of medium- to long-chain unbranched 3-oxo acyl-CoAs into acetyl-CoA and a fatty acyl-CoA shortened by two carbon atoms (Probable). Also catalyzes the condensation of two acetyl-CoA molecules into acetoacetyl-CoA and could be involved in the production of ketone bodies (Probable). Also displays hydrolase activity on various fatty acyl-CoAs (PubMed:25478839). Thereby, could be responsible for the production of acetate in a side reaction to beta-oxidation (Probable). Abolishes BNIP3-mediated apoptosis and mitochondrial damage (PubMed:18371312) In the production of energy from fats, this is one of the enzymes that catalyzes the last step of the mitochondrial beta-oxidation pathway, an aerobic process breaking down fatty acids into acetyl-CoA. Using free coenzyme A/CoA, catalyzes the thiolitic cleavage of medium- to long-chain unbranched 3-oxoacyl-CoAs into acetyl-CoA and a fatty acyl-CoA shortened by two carbon atoms. Also catalyzes the condensation of two acetyl-CoA molecules into acetoacetyl-CoA and could be involved in the production of ketone bodies. Also displays hydrolase activity on various fatty acyl-CoAs (By similarity). Thereby, could be responsible for the production of acetate in a side reaction to beta-oxidation (By similarity). Abolishes BNIP3-mediated apoptosis and mitochondrial damage (By similarity) In the production of energy from fats, this is one of the enzymes that catalyzes the last step of the mitochondrial beta-oxidation pathway, an aerobic process breaking down fatty acids into acetyl-CoA (Probable). Using free coenzyme A/CoA, catalyzes the thiolitic cleavage of medium- to long-chain unbranched 3-oxoacyl-CoAs into acetyl-CoA and a fatty acyl-CoA shortened by two carbon atoms (Probable). Also catalyzes the condensation of two acetyl-CoA molecules into acetoacetyl-CoA and could be involved in the production of ketone bodies (Probable). Also displays hydrolase activity on various fatty acyl-CoAs (PubMed:16476568). Thereby, could be responsible for the production of acetate in a side reaction to beta-oxidation (Probable). Abolishes BNIP3-mediated apoptosis and mitochondrial damage (By similarity)
基因ID	10449
基因名	ACAA2, Acaa2
Swiss	P42765, Q8BWT1, P13437
别名	ACAA2 (YD36015)

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn