— ABLYBIO, Help Your Research

# ALK (YD18645) Rabbit mAb

**货号: AYD16158**

## 产品信息

| | |
|---|---|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC |
| 推荐浓度 | |
| 理论分子量 | 176kDa |
| 实测分子量 | |
| 形式 | Liquid |
| 保存条件 | Store at -20℃. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | PC-3,Mouse testis |
| 细胞定位 | Cell membrane |
| 纯化 | 亲和纯化 |

## 抗原信息

| | |
|---|---|
| 抗原信息 | |

## 靶点信息

| | |
|---|---|
| 研究背景 | This gene encodes a receptor tyrosine kinase, which belongs to the insulin receptor superfamily. This protein comprises an extracellular domain, an hydrophobic stretch corresponding to a single pass transmembrane region, and an intracellular kinase domain. It plays an important role in the development of the brain and exerts its effects on specific neurons in the nervous system. This gene has been found to be rearranged, mutated, or amplified in a series of tumours including anaplastic large cell lymphomas, neuroblastoma, and non-small cell lung cancer. The chromosomal rearrangements are the most common genetic alterations in this gene, which result in creation of multiple fusion genes in tumourigenesis, including ALK (chromosome 2)/EML4 (chromosome 2), ALK/RANBP2 (chromosome 2), ALK/ATIC (chromosome 2), ALK/TFG (chromosome 3), ALK/NPM1 (chromosome 5), ALK/SQSTM1 (chromosome 5), ALK/KIF5B (chromosome 10), ALK/CLTC (chromosome 17), ALK/TPM4 (chromosome 19), and ALK/MSN (chromosome X). |
| 基因ID | 238 |
| 基因名 | ALK |
| Swiss | Q9UM73 (https://www.uniprot.org/uniprotkb/Q9UM73/entry) |
| 别名 | ALK (YD18645),ALK (YD18645) Rabbit mAb,ALK,Anaplastic lymphoma kinase |

## 产品验证

## 实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)