

CDC40 (YD36061) Rabbit mAb

货号: **AYD16296**

产品信息

反应	Human, Mouse, Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P ICC/IF FC
推荐浓度	
理论分子量	66kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HCT116,HepG2,Mouse lung,Mouse brain,Mouse spleen,Rat lung
细胞定位	Nucleus, Nucleus speckle
纯化	

抗原信息


抗原信息	
------	--

靶点信息

研究背景	Pre-mRNA splicing occurs in two sequential transesterification steps. The protein encoded by this gene is found to be essential for the catalytic step II in pre-mRNA splicing process. It is found in the spliceosome, and contains seven WD repeats, which function in protein-protein interactions. This protein has a sequence similarity to yeast Prp17 protein, which functions in two different cellular processes: pre-mRNA splicing and cell cycle progression. It suggests that this protein may play a role in cell cycle progression. [provided by RefSeq, Jul 2008]
基因ID	51362
基因名	CDC40

Swiss	O60508
别名	CDC40 (YD36061)

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn