

— ABLYBIO, Help Your Research



RFC4 (YD36287) Rabbit mAb

货号: **AYD16525**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB ICC/IF IP
推荐浓度	
理论分子量	40kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,SW480,22Rv1,SW620,K-562,HepG2,MCF7
细胞定位	Nucleus
纯化	亲和纯化



抗原信息

抗原信息	
------	--

靶点信息

研究背景	The elongation of primed DNA templates by DNA polymerase delta and DNA polymerase epsilon requires the accessory proteins proliferating cell nuclear antigen (PCNA) and replication factor C (RFC). RFC, also named activator 1, is a protein complex consisting of five distinct subunits of 140, 40, 38, 37, and 36 kD. This gene encodes the 37 kD subunit. This subunit forms a core complex with the 36 and 40 kDa subunits. The core complex possesses DNA-dependent ATPase activity, which was found to be stimulated by PCNA in an in vitro system. Alternatively spliced transcript variants encoding the same protein have been reported.
基因ID	5984
基因名	RFC4
Swiss	P35249 (https://www.uniprot.org/uniprotkb/P35249/entry)
别名	RFC4 (YD36287),RFC4 (YD36287) Rabbit mAb,RFC4,Activator 1 37 kDa subunit,Activator 1 subunit 4,Replication factor C 37 kDa subunit

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)