

# CD16/CD32 (YD35724) Rabbit mAb

货号: **AYD16577**

## 产品信息

反应	Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC-P ICC/IF FC IP
推荐浓度	
理论分子量	37kDa/30kDa/30kDa/30kDa/30kDa/30kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cell membrane, Cytoplasm, cytoskeleton, Secreted
纯化	

## 抗原信息

抗原信息	
------	--

## 靶点信息

研究背景	Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of antigen-antibody complexes from the circulation and modulation of antibody production by B-cells. Isoform IIB1 and isoform IIB1' form caps but fail to mediate endocytosis or phagocytosis. Isoform IIB2 can mediate the endocytosis of soluble immune complexes via clathrin-coated pits. Isoform IIB1 and isoform IIB2 can down-regulate B-cell, T-cell, and mast cell activation when coaggregated to B-cell receptors for AG (BCR), T-cell receptors for AG (TCR), and Fc receptors, respectively Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor which binds to IgG1, IgG2a and IgG2b (PubMed:17558411). Mediates neutrophil activation by IgG complexes redundantly with Fcgr4 (PubMed:18097064)
基因ID	3134

基因名	Fcgr2, Fcgr3
Swiss	P08101; P08508; Q5D5I8; Q5D5J5; Q7TMW9; Q9ES92
别名	CD16/CD32 (YD35724)

产品验证

实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)