

— ABLYBIO, Help Your Research



# TAOK3 (YD35846) Rabbit mAb

货号: **AYD16587**

## 产品信息

反应	Human, Mouse, Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	
理论分子量	105kDa/105kDa/105kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cytoplasm, Cell membrane, Membrane raft, Lipid droplet
纯化	亲和纯化

## 抗原信息

抗原信息	
------	--

## 靶点信息

研究背景	<p>Serine/threonine-protein kinase that acts as a regulator of the p38/MAPK14 stress-activated MAPK cascade and of the MAPK8/JNK cascade. In response to DNA damage, involved in the G2/M transition DNA damage checkpoint by activating the p38/MAPK14 stress-activated MAPK cascade, probably by mediating phosphorylation of upstream MAP2K3 and MAP2K6 kinases. Inhibits basal activity of the MAPK8/JNK cascade and diminishes its activation in response to epidermal growth factor (EGF). Positively regulates canonical T cell receptor (TCR) signaling by preventing early PTPN6/SHP1-mediated inactivation of LCK, ensuring sustained TCR signaling that is required for optimal activation and differentiation of T cells (PubMed:30373850). Phosphorylates PTPN6/SHP1 on 'Thr-394', leading to its polyubiquitination and subsequent proteasomal degradation (PubMed:38166031). Required for cell surface expression of metalloprotease ADAM10 on type 1 transitional B cells which is necessary for their NOTCH-mediated development into marginal zone B cells (By similarity). Also required for the NOTCH-mediated terminal differentiation of splenic conventional type 2 dendritic cells (By similarity). Positively regulates osteoblast differentiation by acting as an upstream activator of the JNK pathway (PubMed:32807497). Promotes JNK signaling in hepatocytes and positively regulates hepatocyte lipid storage by inhibiting beta-oxidation and triacylglycerol secretion while enhancing lipid synthesis (PubMed:34634521). Restricts age-associated inflammation by negatively regulating differentiation of macrophages and their production of pro-inflammatory cytokines (By similarity). Plays a role in negatively regulating the abundance of regulatory T cells in white adipose tissue (By similarity)</p> <p>Serine/threonine-protein kinase that acts as a regulator of the p38/MAPK14 stress-activated MAPK cascade and of the MAPK8/JNK cascade (By similarity). In response to DNA damage, involved in the G2/M transition DNA damage checkpoint by activating the p38/MAPK14 stress-activated MAPK cascade, probably by mediating phosphorylation of upstream MAP2K3 and MAP2K6 kinases (By similarity). Inhibits basal activity of the MAPK8/JNK cascade and diminishes its activation in response to epidermal growth factor (EGF) (By similarity). Positively regulates canonical T cell receptor (TCR) signaling by preventing early PTPN6/SHP1-mediated inactivation of LCK, ensuring sustained TCR signaling that is required for optimal activation and differentiation of T cells (By similarity). Phosphorylates PTPN6/SHP1 on 'Thr-394', leading to its polyubiquitination and subsequent proteasomal degradation (By similarity). Required for cell surface expression of metalloprotease ADAM10 on type 1 transitional B cells which is necessary for their NOTCH-mediated development into marginal zone B cells (PubMed:28068307). Also required for the NOTCH-mediated terminal differentiation of splenic conventional type 2 dendritic cells (PubMed:33214146). Positively regulates osteoblast differentiation by acting as an upstream activator of the JNK pathway (PubMed:32807497). Promotes JNK signaling in hepatocytes and positively regulates hepatocyte lipid storage by inhibiting beta-oxidation and triacylglycerol secretion while enhancing lipid synthesis (By similarity). Restricts age-associated inflammation by negatively regulating differentiation of macrophages and their production of pro-inflammatory cytokines (PubMed:37400834). Plays a role in negatively regulating the abundance of regulatory T cells in white adipose tissue (PubMed:37347461)</p> <p>Serine/threonine-protein kinase that acts as a regulator of the p38/MAPK14 stress-activated MAPK cascade and of the MAPK8/JNK cascade. In response to DNA damage, involved in the G2/M transition DNA damage checkpoint by activating the p38/MAPK14 stress-activated MAPK cascade, probably by mediating phosphorylation of upstream MAP2K3 and MAP2K6 kinases. Inhibits basal activity of the MAPK8/JNK cascade and diminishes its activation in response to epidermal growth factor (EGF). Positively regulates canonical T cell receptor (TCR) signaling by preventing early PTPN6/SHP1-mediated inactivation of LCK, ensuring sustained TCR signaling that is required for optimal activation and differentiation of T cells. Phosphorylates PTPN6/SHP1 on 'Thr-396', leading to its polyubiquitination and subsequent proteasomal degradation. Required for cell surface expression of metalloprotease ADAM10 on type 1 transitional B cells which is necessary for their NOTCH-mediated development into marginal zone B cells. Also required for the NOTCH-mediated terminal differentiation of splenic conventional type 2 dendritic cells. Positively regulates osteoblast differentiation by acting as an upstream activator of the JNK pathway. Promotes JNK signaling in hepatocytes and positively regulates hepatocyte lipid storage by inhibiting beta-oxidation and triacylglycerol secretion while enhancing lipid synthesis. Restricts age-associated inflammation by negatively regulating differentiation of macrophages and their production of pro-inflammatory cytokines. Plays a role in negatively regulating the abundance of regulatory T cells in white adipose tissue</p>
基因ID	51347
基因名	TAOK3, Taok3
Swiss	Q9HC79 ( <a href="https://www.uniprot.org/uniprotkb/Q9HC79/entry">https://www.uniprot.org/uniprotkb/Q9HC79/entry</a> ), Q8BYC6 ( <a href="https://www.uniprot.org/uniprotkb/Q8BYC6/entry">https://www.uniprot.org/uniprotkb/Q8BYC6/entry</a> ), Q53UA7 ( <a href="https://www.uniprot.org/uniprotkb/Q53UA7/entry">https://www.uniprot.org/uniprotkb/Q53UA7/entry</a> )

别名	TAOK3 (YD35846),TAOK3 (YD35846) Rabbit mAb,TAOK3,Cutaneous T-cell lymphoma-associated antigen H D-CL-09,Dendritic cell-derived protein kinase,JNK/SAPK-inhibitory kinase,Jun kinase-inhibitory kinase,Kinase from chicken homolog A,Thousand and one amino acid protein 3
----	---

### 产品验证

### 实验步骤

访问官网浏览详情: [www.ablybio.cn \(https://www.ablybio.cn/www.ablybio.cn\)](https://www.ablybio.cn/www.ablybio.cn)