

— ABLYBIO, Help Your Research



mtTFA (YD34889) Rabbit mAb

货号: **AYD16614**

产品信息

反应	Mouse, Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB ICC/IF FC IP
推荐浓度	
理论分子量	28kDa/28kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Mitochondrion, Mitochondrion matrix, mitochondrion nucleoid, Nucleus
纯化	亲和纯化


抗原信息

抗原信息	
------	--

靶点信息

研究背景	Binds to the mitochondrial light strand promoter and functions in mitochondrial transcription regulation (By similarity). Component of the mitochondrial transcription initiation complex, composed at least of TFB2M, TFAM and POLRMT that is required for basal transcription of mitochondrial DNA (By similarity). In this complex, TFAM recruits POLRMT to a specific promoter whereas TFB2M induces structural changes in POLRMT to enable promoter opening and trapping of the DNA non-template strand (By similarity). Required for accurate and efficient promoter recognition by the mitochondrial RNA polymerase (By similarity). Promotes transcription initiation from the HSP1 and the light strand promoter by binding immediately upstream of transcriptional start sites (By similarity). Is able to unwind DNA (By similarity). Bends the mitochondrial light strand promoter DNA into a U-turn shape via its HMG boxes (By similarity). Required for maintenance of normal levels of mitochondrial DNA (PubMed:9500544). May play a role in organizing and compacting mitochondrial DNA (PubMed:17581862) Binds to the mitochondrial light strand promoter and functions in mitochondrial transcription regulation. Component of the mitochondrial transcription initiation complex, composed at least of TFB2M, TFAM and POLRMT that is required for basal transcription of mitochondrial DNA. In this complex, TFAM recruits POLRMT to a specific promoter whereas TFB2M induces structural changes in POLRMT to enable promoter opening and trapping of the DNA non-template strand. Required for accurate and efficient promoter recognition by the mitochondrial RNA polymerase. Promotes transcription initiation from the HSP1 and the light strand promoter by binding immediately upstream of transcriptional start sites. Is able to unwind DNA. Bends the mitochondrial light strand promoter DNA into a U-turn shape via its HMG boxes. Required for maintenance of normal levels of mitochondrial DNA. May play a role in organizing and compacting mitochondrial DNA (By similarity)
基因ID	3134
基因名	Tfam
Swiss	P40630 (https://www.uniprot.org/uniprotkb/P40630/entry), Q91ZW1 (https://www.uniprot.org/uniprotkb/Q91ZW1/entry)
别名	mtTFA (YD34889),mtTFA (YD34889) Rabbit mAb,Tfam,Testis-specific high mobility group protein,Hmgts

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)