

# FOXJ1 (YD15819) Rabbit mAb

货号: **AYD16686**

## 产品信息

反应	Human, Mouse, Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	IHC-P
推荐浓度	
理论分子量	45kDa/45kDa/46kDa
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Nucleus
纯化	

## 抗原信息

抗原信息	
------	--

## 靶点信息

研究背景	Transcription factor specifically required for the formation of motile cilia (PubMed:31630787). Acts by activating transcription of genes that mediate assembly of motile cilia, such as CFAP157. Binds the DNA consensus sequences 5'-HWDTGTTTGTTA-3' or 5'-KTTTGTTGTTKW-3' (where H is not G, W is A or T, D is not C, and K is G or T). Activates the transcription of a variety of ciliary proteins in the developing brain and lung Transcription factor specifically required for the formation of motile cilia (PubMed:10873152, PubMed:14996907, PubMed:22357932, PubMed:27965440, PubMed:9096351, PubMed:9739041). Acts by activating transcription of genes that mediate assembly of motile cilia, such as CFAP157 (PubMed:27965440). Binds the DNA consensus sequences 5'-HWDTGTTTGTTA-3' or 5'-KTTTGTTGTTKW-3' (where H is not G, W is A or T, D is not C, and K is G or T) (PubMed:9096351). Activates the transcription of a variety of ciliary proteins in the developing brain and lung (PubMed:27914912, PubMed:28666954) Transcription factor specifically required for the formation of motile cilia. Acts by activating transcription of genes that mediate assembly of motile cilia, such as CFAP157. Binds the DNA consensus sequences 5'-HWDTGTTTGTTA-3' or 5'-KTTTGTTGTTKW-3' (where H is not G, W is A or T, D is not C, and K is G or T). Activates the transcription of a variety of ciliary proteins in the developing brain and lung
基因ID	2302
基因名	FOXJ1, Foxj1
Swiss	Q92949, Q61660, Q63247
别名	FOXJ1 (YD15819)

#### 产品验证

#### 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)