

HDAC11 Rabbit mAb

货号: **AYM28404**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	33kDa/39kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,U-87MG,Mouse spinal cord,Rat brain
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human HDAC11.
序列	GLSISPAGIVKRDELVFRMVRGRRVPILMVTSGGYQKRTARIADSILNLFGLGLIGPESPSVSAQNSDTPLLPVAVP

靶点信息

研究背景	This gene encodes a class IV histone deacetylase. The encoded protein is localized to the nucleus and may be involved in regulating the expression of interleukin 10. Alternative splicing results in multiple transcript variants.
基因ID	79885
基因名	HDAC11

Swiss	Q96DB2
别名	HDAC11;HD11

产品验证



Western blot analysis of HDAC11 expressed in MCF7,U-87MG,Mouse spinal cord using HDAC11 Rabbit m Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s

实验步骤

访问官网浏览详情: www.ablybio.cn