

Flotillin 2 Rabbit mAb

货号: **AYM28413**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	47kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,SKOV3,Mouse heart,Mouse lung,Rat brain
细胞定位	Cell membrane,Endosome,Lipid-anchor,Membrane,Peripheral membrane protein,caveola
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Flotillin 2.
序列	LAYELQGAREQQKIRQEEIEIEVVQRKKQIAVEAQEILRTDKELIATVRRPAEAEAHRIQQIAEGEKVKQVLLAQAEAEKIRKIGEAEAAVIEAMGKAEAERMKLKAEAYQKYGDAAKMALVLEALPQIAAKIAAPLTKVDEIVVLSGDNSKVTSEVNRLLAELPASVHALTGVDLSKIPLIKKATGVQV

靶点信息

研究背景	Caveolae are small domains on the inner cell membrane involved in vesicular trafficking and signal transduction. This gene encodes a caveolae-associated, integral membrane protein, which is thought to function in neuronal signaling.
基因ID	2319

基因名	FLOT2
Swiss	Q14254
别名	FLOT2;ECS-1;ECS1;ESA;ESA1;M17S1

产品验证



Western blot analysis of Flotillin 2 expressed in Mouse heart,Rat brain,A-549 using Flotillin 2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 S.

实验步骤

访问官网浏览详情: www.ablybio.cn