

# delta 1 Catenin/p120 Catenin Rabbit mAb

货号: **AYM28431**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	105kDa
实测分子量	105kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,U-87MG,Mouse testis,Rat brain
细胞定位	Cell membrane,Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human delta 1 Catenin/p120 Catenin.
------	---

## 靶点信息

研究背景	This gene encodes a member of the Armadillo protein family, which function in adhesion between cells and signal transduction. Multiple translation initiation codons and alternative splicing result in many different isoforms being translated. Not all of the full-length nature of the described transcript variants have been determined. Read-through transcription also exists between this gene and the neighboring upstream thiodoxin-related transmembrane protein 2 (TMX2) gene. [provided by RefSeq, Dec 2010]
基因ID	1500

基因名	CTNND1
Swiss	O60716
别名	delta 1 Catenin/p120 Catenin

## 产品验证



Western blot analysis of delta 1 Catenin/p120 Catenin expressed in Mouse testis,Rat brain,HeLa using delta 1 Catenin/p120 Catenin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)