

— ABLYBIO, Help Your Research



SIRT1 Rabbit mAb

货号: **AYM28446**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP IF
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	61kDa/81kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cytoplasm,Mitochondrion,Nucleus,Nucleus,PML body
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human SIRT1.
------	--


靶点信息

研究背景	This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Alternative splicing results in multiple transcript variants.
基因ID	23411
基因名	SIRT1
Swiss	Q96EB6 (https://www.uniprot.org/uniprotkb/Q96EB6/entry)
别名	SIRT1, 75SirT1, hSIR2, hSIRT1, HST2, S. cerevisiae, homolog of, NAD dependent deacetylase sirtuin 1, NAD dependent protein deacetylase sirtuin 1, OTTHUMP00000198111, OTTHUMP00000198112, Regulatory protein SIR2 homolog 1, SIR1_HUMAN, SIR2, SIR2 like 1, SIRT1 Rabbit mAb, NAD-dependent protein deacetylase sirtuin -1, SIR2-like protein 1, SIR2L1


产品验证




Western blot analysis of SIRT1 expressed in 293T using SIRT1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.




Immunohistochemical analysis of paraffin-embedded Rat stomach, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Human pituitary tumor, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Human colon, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Human placenta, using the Antibody at 1:50 dilution.



Immunohistochemical analysis of paraffin-embedded Mouse lung, using the Antibody at 1:50 dilution.



Immunohistochemical analysis of paraffin-embedded human kidney, using SIRT1 Antibody.



Immunofluorescence analysis of SIRT1 cells labelling AKT1/2/3 with . The cells were fixed with cold 100% methanol (10min, 4°C) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with Goat Anti-Rabbit IgG Fluor488 (1:200, shown in green) at room temperature for 1h.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)