

Survivin Rabbit mAb

货号: **AYM28457**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	8kDa/13kDa/15kDa/16kDa/18kDa
实测分子量	16kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293F,Jurkat
细胞定位	Chromosome,Cytoplasm,Midbody,Nucleus,centromere,cytoskeleton,kinetochore,spindle
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Survivin.
序列	MGAPTLPPAWQPFLKDHRIKSTFKNWPFLGCACTPERMAEAGFIHCPTENEPDLAQCFKCFKELEGWEPDDDDPIEEHKKH SSGCAFLSVKKQFEELTLGFLKLDREKAKNKIAKETNNKKKEFEETAEKVRRRAIEQLAAMD


靶点信息

研究背景	This gene is a member of the inhibitor of apoptosis (IAP) gene family, which encode negative regulatory proteins that prevent apoptotic cell death. IAP family members usually contain multiple baculovirus IAP repeat (BIR) domains, but this gene encodes proteins with only a single BIR domain. The encoded proteins also lack a C-terminus RING finger domain. Gene expression is high during fetal development and in most tumors, yet low in adult tissues. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.
基因ID	332
基因名	BIRC5
Swiss	O15392
别名	API4;EPR-1;BIRC5;Survivin

产品验证



Immunohistochemical analysis of paraffin-embedded human gastric carcinoma, using Survivin Antibody.



Western blot analysis of Survivin expressed in HeLa, 293F, Jurkat using Survivin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn