

— ABLYBIO, Help Your Research



CEBP Beta Rabbit mAb

货号: **AYM28474**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC IP
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50 IP: 1:20 - 1:50
理论分子量	15kDa/33kDa/36kDa
实测分子量	33kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SK-MEL-28,HeLa,NIH/3T3
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CEBP Beta.
------	--

靶点信息

研究背景	This intronless gene encodes a transcription factor that contains a basic leucine zipper (bZIP) domain. The encoded protein functions as a homodimer but can also form heterodimers with CCAAT/enhancer-binding proteins alpha, delta, and gamma. Activity of this protein is important in the regulation of genes involved in immune and inflammatory responses, among other processes. The use of alternative in-frame AUG start codons results in multiple protein isoforms, each with distinct biological functions.
基因ID	1051
基因名	CEBPB
Swiss	P17676 (https://www.uniprot.org/uniprotkb/P17676/entry)
别名	CEBP Beta,CEBP Beta Rabbit mAb,CEBPB,Liver activator protein,Liver-enriched inhibitory protein,Nuclear factor NF-IL6,Transcription factor 5,TCF5

产品验证



Western blot analysis of CEBP Beta expressed in SK-MEL-28, HeLa, NIH/3T3 using CEBP Beta Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)