

# NFIB / NF1B2 Rabbit mAb

货号: **AYM28503**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50 <b>FC:</b> 1:20 - 1:50
理论分子量	22kDa/33kDa/47kDa/53kDa/55kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human NFIB / NF1B2.
序列	SRTWHERDQDMSSPTTMKKPEKPLFSSASPQDSSPRLSTFPQHHPGIPGVAHSVISTRTPPPSPLPFPTQAILPPAPSSYFSHTIRYPPHLNPQDTLKNYVPSYDPSSPQTSQSWYLG

## 靶点信息

研究背景	Transcriptional activator of GFAP, essential for proper brain development. Recognizes and binds the palindromic sequence 5'-TTGGCNNNNNGCCAA-3' present in viral and cellular promoters and in the origin of replication of adenovirus type 2. These proteins are individually capable of activating transcription and replication.
------	--

基因ID	4781
基因名	NFIB
Swiss	O00712
别名	NFIB;CTF;HMGIC/NFIB;NF-I/B;NF1-B;NFI-B;NFI-RED;NFIB2;NFIB3

## 产品验证



Western blot analysis of NFIB / NF1B2 expressed in HeLa using NFIB / NF1B2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)