

htrA1 Rabbit mAb

货号: **AYM28505**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	51kDa
实测分子量	51kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Mouse liver,Rat brain
细胞定位	Cytoplasm,Secreted,cytosol
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human htrA1.
序列	AQLSRAGRSAPLAAGCPDRCEPARCPPQPEHCEGGRARDACGCCEVCGAPEGAACGLQEGPCGEGLCVVPFGVPASATVRRRAQAGLCVCASSEPVCGSDANTYANLCQLRAASRRSERLHRPPVIVLQRGACGQGQE

靶点信息

研究背景	This gene encodes a member of the trypsin family of serine proteases. This protein is a secreted enzyme that is proposed to regulate the availability of insulin-like growth factors (IGFs) by cleaving IGF-binding proteins. It has also been suggested to be a regulator of cell growth. Variations in the promoter region of this gene are the cause of susceptibility to age-related macular degeneration type 7.
基因ID	5654

基因名	HTRA1
Swiss	Q92743
别名	HTRA1;ARMD7;CADASIL2;CARASIL;HtrA;L56;ORF480;PRSS11

产品验证



Western blot analysis of htrA1 expressed in Mouse brain, Mouse liver, Rat brain using htrA1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn